

Overdispersion

Recap: Overdispersion

Overdispersion occurs when the response Y has higher variance than we would expect from the specified EDM

$$\text{var}(\hat{\beta} \mid \phi > 1) = \phi \text{var}(\hat{\beta} \mid \phi = 1)$$

Poisson regression: $\phi = 1$ (we use this to estimate $\text{var}(\hat{\beta})$)

$\Rightarrow$ overdispersion means estimated variances & SEs are too small

Estimating ϕ

Intuition: $Y_i \sim N(\mu_i, \sigma^2)$ ↙ dispersion

$$\hat{\sigma}^2 = \frac{SSE}{n-p} = \frac{SSE}{n-(k+1)} = \frac{1}{n-(k+1)} \sum_{i=1}^n (Y_i - \hat{Y}_i)^2$$

How to generalize this?

Option: Recall $D(y, \mu) = (y - \mu)^2$
 $\Rightarrow D(y, \hat{\mu}) = \sum_{i=1}^n (Y_i - \hat{Y}_i)^2 = SSE$

$$\text{So } \hat{\sigma}^2 = \frac{D(y, \hat{\mu})}{n-(k+1)}$$

Generalize this:

$$\hat{\phi} = \frac{D(y, \hat{\mu})}{n-(k+1)}$$

(mean variance estimate)

Saddlepoint approximation

$$f(y; \mu, \phi) \approx \frac{1}{\sqrt{2\pi\phi v(y)}} \exp \left\{ -\frac{\delta(y, \mu)}{2\phi} \right\}$$

$$\text{Log likelihood: } \ell(\mu, \phi) \approx \sum_{i=1}^n \left\{ -\frac{1}{2} \log(2\pi\phi v(y_i)) - \frac{1}{2\phi} \delta(y_i, \mu_i) \right\}$$

$$\frac{\partial \ell(\mu, \phi)}{\partial \phi} = \sum_{i=1}^n \left\{ -\frac{1}{2} \cdot \frac{1}{\phi} + \frac{1}{2\phi^2} \delta(y_i, \mu_i) \right\} \stackrel{\text{set}}{=} 0$$

$$\Rightarrow \sum_{i=1}^n -\frac{1}{2} \cdot \frac{1}{\phi} + \frac{1}{2\phi^2} \sum_{i=1}^n \delta(y_i, \mu_i) = 0$$

$$\Rightarrow -\frac{n}{2} \cdot \frac{1}{\phi} + \frac{1}{2\phi^2} \sum_{i=1}^n \delta(y_i, \mu_i) = 0$$

$$\Rightarrow \hat{\phi}_{MLE} = \frac{1}{n} \sum_{i=1}^n \delta(y_i, \mu_i) = \frac{D(y, \mu)}{n}$$

we don't know μ

$$\Rightarrow \hat{\phi}_{MLE} = \frac{D(y, \hat{\mu})}{n} \quad (\text{plug in } \hat{\mu})$$

But, this is biased

As usual with variance estimators:

$$\hat{\phi} = \frac{D(y, \hat{\mu})}{n - (k+1)}$$

mean variance
estimate

$$\hat{\sigma}^2 = \frac{1}{n - (k+1)} \sum_{i=1}^n (y_i - \hat{y}_i)^2 \quad v(\mu_i) = 1$$

Option 2:
$$\sum_{i=1}^n (y_i - \hat{y}_i)^2 = \sum_{i=1}^n \frac{(y_i - \hat{y}_i)^2}{1} = \sum_{i=1}^n \frac{(y_i - \hat{\mu}_i)^2}{v(\hat{\mu}_i)}$$

Pearson residuals:
$$\frac{y_i - \hat{\mu}_i}{\sqrt{v(\hat{\mu}_i)}}$$

intuition: account for mean-variance relationship in residuals

$$\Rightarrow \hat{\phi} = \frac{1}{n - (k+1)} \sum_{i=1}^n \frac{(y_i - \hat{\mu}_i)^2}{v(\hat{\mu}_i)}$$

← sum of squared Pearson residuals

↑
Pearson estimate

Ex: Poisson regression

$$(v(\mu_i) = \mu_i)$$

$$\text{Pearson estimate} = \frac{1}{n - (k+1)} \sum_{i=1}^n \frac{(y_i - \hat{\lambda}_i)^2}{\hat{\lambda}_i}$$

Motivation: $\text{var}(y_i) = \phi v(\mu_i)$
Suppose μ_i is known $\Rightarrow \hat{\phi} = \frac{1}{n} \sum_{i=1}^n \frac{(y_i - \mu_i)^2}{v(\mu_i)}$

$$E[\hat{\phi}] = E\left[\frac{(y_i - \mu_i)^2}{v(\mu_i)}\right] = \frac{\text{var}(y_i)}{v(\mu_i)} = \phi$$

which estimate $\hat{\theta}$ should we use?

- the mean-variance estimate depends on the saddlepoint approximation
- the Pearson estimate is approximately unbiased if μ & $v(\mu)$ are correct — we just need the first two moments

$\Rightarrow$ Pearson estimate is more "robust"

- R typically uses the Pearson estimate

Using $\hat{\phi}$

```
pearson_resids <- residuals(m2, type="pearson")
sum(pearson_resids^2)/df.residual(m2)
```

```
## [1] 7.58542
```

$\hat{\phi} = 7.59$ (Pearson estimate)
 $\gg 1$

...

	Estimate	Std. Error	z value	Pr(> z)
## (Intercept)	-1.30445	0.12403	-10.517	< 2e-16 ***
## regionMW	0.09754	0.17752	0.549	0.58270
## regionNE	0.76268	0.15292	4.987	6.12e-07 ***
## regionSE	0.87237	0.15313	5.697	1.22e-08 ***
## regionSW	0.50708	0.18507	2.740	0.00615 **
## regionW	0.20934	0.18605	1.125	0.26053

...

CI for β_S :

$$\underbrace{0.209}_{\hat{\beta}_S} \pm z_{1-\frac{\alpha}{2}} \underbrace{(\sqrt{7.59})}_{\sqrt{\hat{\phi}}} (0.186)$$

Quasi-Poisson regression

A model for overdispersed Poisson-like counts, using an estimated dispersion parameter $\hat{\phi}$, is called a *quasi-Poisson* model.

```
m3 <- glm(nv ~ region, offset = log(enroll1000),
          data = crimes, family = quasipoisson)
summary(m3)
```

...

##		Estimate	Std. Error	t value	Pr(> t)	
##	(Intercept)	-1.30445	0.34161	-3.818	0.000274	***
##	regionMW	0.09754	0.48893	0.199	0.842417	
##	regionNE	0.76268	0.42117	1.811	0.074167	.
##	regionSE	0.87237	0.42175	2.068	0.042044	*
##	regionSW	0.50708	0.50973	0.995	0.323027	

...

↙ already include $\sqrt{\hat{\phi}}$

$$\hat{\phi} \approx \chi^2$$

using canonical link:

$$u(\beta) = \frac{X^T(Y - \mu)}{\phi} \stackrel{\text{set}}{=} 0$$

Poisson vs. quasi-Poisson

Poisson:

Solving $X^T(Y - \mu) = 0$
Wald test (assume $\phi=1$)

...					
##	Estimate	Std. Error	z value	Pr(> z)	
## (Intercept)	-1.30445	0.12403	-10.517	< 2e-16	***
## regionMW	0.09754	0.17752	0.549	0.58270	
## regionNE	0.76268	0.15292	4.987	6.12e-07	***
## regionSE	0.87237	0.15313	5.697	1.22e-08	***
...					

Quasi-Poisson:

↑
same β s

$$SE(\text{quasi Poisson}) = \sqrt{\hat{\phi}} SE(\text{Poisson})$$

e.g., $0.421 = \sqrt{7.59} (0.153)$

...					
##	Estimate	Std. Error	t value	Pr(> t)	$\frac{\beta - 0}{SE(\hat{\beta})}$
## (Intercept)	-1.30445	0.34161	-3.818	0.000274	***
## regionMW	0.09754	0.48893	0.199	0.842417	
## regionNE	0.76268	0.42117	1.811	0.074167	.

increases

p-values 6/7

Quasi-likelihood models

what is the Quasi-Poisson model?

• $V(\mu_i) = \mu_i$ $\text{Var}(Y_i) = \phi \mu_i$

• mean-variance relationship uniquely determines EDM

$\Rightarrow$ the only EDM with $V(\mu_i) = \mu_i$
is a Poisson

Poisson: $\phi = 1$

$\Rightarrow$ quasi-Poisson doesn't correspond to an EDM