


ZIP models

- Challenge 8 posted on course website
- Project 2 released later today
- Hw 6 due Monday
- Exam 2 released next Friday
 - Poisson, EDMs, quasi-Poisson, negative binomial, ZIP models



$y_i =$ bill length

$$y_i = \beta_0 + \beta_1 \text{Mass}_i + \epsilon_i$$

- = Adélie
- = Gentoo
- = Chinstrap

$$y_i = \beta_0 + \beta_1 \text{Mass}_i + \beta_2 \text{Chinstrap}_i + \beta_3 \text{Gentoo}_i + \epsilon_i$$

For Adélie:


$$y_i = \beta_0 + \beta_1 \text{Mass}_i + \epsilon_i$$


Chinstrap:

$$y_i = (\beta_0 + \beta_2) + \beta_1 \text{Mass}_i + \epsilon_i$$

Gentoo:

$$y_i = (\beta_0 + \beta_3) + \beta_1 \text{Mass} + \epsilon_i$$





- = Adelie
- = Gentoo
- = Chinstrap

$$\hat{y}_i = \beta_0 + \beta_1 \text{Mass}_i + \beta_2 \text{Chinstrap}_i + \beta_3 \text{Gentoo}_i +$$

$$\beta_4 \cdot \text{Mass}_i \cdot \text{Chinstrap}_i + \beta_5 \cdot \text{Mass}_i \cdot \text{Gentoo}_i + \epsilon_i$$

β_1 = slope for Adelies

β_4 = change in slope for Chinstrap vs. Adelies

β_5 = change in slope for Gentoo vs. Adelies

H_0 : all species have same slope
 $\beta_4 = \beta_5 = 0$

For Adelie:

$$\hat{y}_i = \beta_0 + \beta_1 \text{Mass}_i + \epsilon_i$$

Chinstrap:

$$\hat{y}_i = (\beta_0 + \beta_2) + (\beta_1 + \beta_4) \text{Mass}_i + \epsilon_i$$

Gentoo:

$$\hat{y}_i = (\beta_0 + \beta_3) + (\beta_1 + \beta_5) \text{Mass}_i + \epsilon_i$$

Recap: Zero-inflated Poisson (ZIP) model

$$P(Y_i = y) = \begin{cases} e^{-\lambda_i}(1 - \alpha_i) + \alpha_i & y = 0 \\ \frac{e^{-\lambda_i} \lambda_i^y}{y!} (1 - \alpha_i) & y > 0 \end{cases}$$

mixture of a point mass at 0 and a Poisson
 α_i = mixing proportion

where
logistic component

$$\log\left(\frac{\alpha_i}{1 - \alpha_i}\right) = \gamma_0 + \gamma_1 \text{FirstYear}_i + \gamma_2 \text{OffCampus}_i + \gamma_3 \text{Male}_i$$

$$\log(\lambda_i) = \beta_0 + \beta_1 \text{FirstYear}_i + \beta_2 \text{OffCampus}_i + \beta_3 \text{Male}_i$$

Poisson component

$$\alpha_i = P(Z_i = 1)$$

$$Y_i | (Z_i = 1) \equiv 0 \quad (\text{point mass})$$

$$Y_i | (Z_i = 0) \sim \text{Poisson}(\lambda_i)$$

Fitted model

$$P(Y_i = y) = \begin{cases} e^{-\lambda_i}(1 - \alpha_i) + \alpha_i & y = 0 \\ \frac{e^{-\lambda_i}\lambda_i^y}{y!}(1 - \alpha_i) & y > 0 \end{cases}$$

$$\log\left(\frac{\hat{\alpha}_i}{1 - \hat{\alpha}_i}\right) = -0.40 + 0.89FirstYear_i - 1.69OffCampus_i - 0.07Male_i$$

$$\log(\hat{\lambda}_i) = 0.80 - 0.16FirstYear_i + 0.37OffCampus_i + 0.98Male_i$$

Warm up: Class activity

https://sta712-f22.github.io/class_activities/ca_lecture_29.html

Class activity

$\lambda_i = \# \text{ drinks consumed}$

$Z_i = \mathbb{1}\{\text{student } i \text{ never drinks}\}$

$$P(Y_i = y) = \begin{cases} e^{-\lambda_i}(1 - \alpha_i) + \alpha_i & y = 0 \\ \frac{e^{-\lambda_i} \lambda_i^y}{y!} (1 - \alpha_i) & y > 0 \end{cases}$$

$$\log\left(\frac{\hat{\alpha}_i}{1 - \hat{\alpha}_i}\right) = \underbrace{-0.40 + 0.89\text{FirstYear}_i - 1.69\text{OffCampus}_i - 0.07\text{Male}_i}_{\log(\hat{\lambda}_i)}$$

$$\log(\hat{\lambda}_i) = 0.80 - 0.16\text{FirstYear}_i + 0.37\text{OffCampus}_i + 0.98\text{Male}_i$$

What is the estimated probability that a male first year student who lives on campus *never* drinks?

$$\begin{aligned} P(Z_i = 1) &= \alpha_i \\ \hat{\alpha}_i &= \frac{\exp\{-0.4 + 0.89 - 0.07\}}{1 + \exp\{-0.4 + 0.89 - 0.07\}} = 0.60 \end{aligned}$$

Class activity

$$P(Y_i = y) = \begin{cases} e^{-\lambda_i}(1 - \alpha_i) + \alpha_i & y = 0 \\ \frac{e^{-\lambda_i} \lambda_i^y}{y!} (1 - \alpha_i) & y > 0 \end{cases}$$

$$\log\left(\frac{\hat{\alpha}_i}{1 - \hat{\alpha}_i}\right) = -0.40 + 0.89\text{FirstYear}_i - 1.69\text{OffCampus}_i - 0.07\text{Male}_i$$

$$\log(\hat{\lambda}_i) = 0.80 - 0.16\text{FirstYear}_i + 0.37\text{OffCampus}_i + 0.98\text{Male}_i$$

What is the estimated probability that a male first year student who lives on campus consumed 3 drinks last weekend?

$$\hat{P}(Y_i = 3) = \frac{e^{-\hat{\lambda}_i} \hat{\lambda}_i^3}{3!} (1 - \hat{\alpha}_i) \quad \hat{\alpha}_i = 0.60$$

$$\hat{\lambda}_i = \exp\{0.8 - 0.16 + 0.98\} = 5.05$$

$$\hat{P}(Y_i = 3) = 0.055$$

$$P(Y_i, Z_i | \gamma, \beta) = \underbrace{P(Z_i | \gamma)}_{P(Z_i | \gamma)} \underbrace{P(Y_i | Z_i, \gamma, \beta)}_{P(Y_i | Z_i, \beta)} \quad P(A \cap B) = P(A)P(B|A)$$

Fitting ZIP models

Suppose we can actually observe latent variable Z_i
 $Z_i \sim \text{Bernoulli}(\alpha_i)$ $Y_i | (Z_i=0) \sim \text{Poisson}(\lambda_i)$

$$\log\left(\frac{\alpha_i}{1-\alpha_i}\right) = \gamma^T X_i$$

$$\log(\lambda_i) = \beta^T X_i$$

$$L(\gamma, \beta) = \prod_{i=1}^n P(Y_i, Z_i | \gamma, \beta) = \prod_{i=1}^n P(Z_i | \gamma) P(Y_i | Z_i, \beta)$$

$$= \prod_{i=1}^n \alpha_i^{Z_i} (1-\alpha_i)^{1-Z_i} \left(\frac{e^{-\lambda_i} \lambda_i^{Y_i}}{Y_i!} \right)^{1-Z_i}$$

$$\Rightarrow \ell(\gamma, \beta) = \sum_{i=1}^n (Z_i \log(\alpha_i) + (1-Z_i) \log(1-\alpha_i)) + \sum_{i=1}^n (1-Z_i) (-\lambda_i + Y_i \log \lambda_i) - \sum_{i=1}^n (1-Z_i) \log(Y_i!)$$

$$\ell(\gamma) = \sum_{i=1}^n (Z_i \log(\alpha_i) + (1-Z_i) \log(1-\alpha_i))$$

$$\ell(\beta) = \sum_{i=1}^n (1-Z_i) (-\lambda_i + Y_i \log \lambda_i)$$

if we know Z_i ,
 I can separately
 maximize $\ell(\gamma)$ and
 $\ell(\beta)$